

Fifth annual conference  
of the SFB-TRR 195

## Towards the Classification of Symplectic Linear Quotient Singularities Admitting a Symplectic Resolution

Johannes Schmitt  
TU Kaiserslautern  
16th September 2021

Joint work with  
Gwyn Bellamy and Ulrich Thiel  
Math. Z. (2021), to appear

# 1. The Classification Problem

2. The Groups in Question

3. Symplectic Reflection Algebras

4. Conclusion

# Symplectic Groups

Let  $V$  be a finite dimensional  $\mathbb{C}$ -vector space.

# Symplectic Groups

Let  $V$  be a finite dimensional  $\mathbb{C}$ -vector space.

The space  $V$  is called **symplectic** if there is a bilinear form  $\omega : V \times V \rightarrow \mathbb{C}$  which is non-degenerate and alternating, i.e.  $\omega(v, v) = 0$  for all  $v \in V$ .

# Symplectic Groups

Let  $V$  be a finite dimensional  $\mathbb{C}$ -vector space.

The space  $V$  is called **symplectic** if there is a bilinear form  $\omega : V \times V \rightarrow \mathbb{C}$  which is non-degenerate and alternating, i.e.  $\omega(v, v) = 0$  for all  $v \in V$ .

For  $V$  symplectic define

$$\mathrm{Sp}(V) := \{g \in \mathrm{GL}(V) \mid \omega(gv, gw) = \omega(v, w)\} \leq \mathrm{GL}(V) .$$

# Symplectic Groups

Let  $V$  be a finite dimensional  $\mathbb{C}$ -vector space.

The space  $V$  is called **symplectic** if there is a bilinear form  $\omega : V \times V \rightarrow \mathbb{C}$  which is non-degenerate and alternating, i.e.  $\omega(v, v) = 0$  for all  $v \in V$ .

For  $V$  symplectic define

$$\mathrm{Sp}(V) := \{g \in \mathrm{GL}(V) \mid \omega(gv, gw) = \omega(v, w)\} \leq \mathrm{GL}(V) .$$

## Example

For  $V = \mathbb{C}^{2n}$  and  $\omega(v, w) := v^\top J_n w$  with  $J_n := \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$ , we have

$$\mathrm{Sp}_{2n}(\mathbb{C}) = \{g \in \mathrm{GL}_{2n}(\mathbb{C}) \mid g^\top J_n g = J_n\} .$$

# Symplectic Groups

Let  $V$  be a finite dimensional  $\mathbb{C}$ -vector space.

The space  $V$  is called **symplectic** if there is a bilinear form  $\omega : V \times V \rightarrow \mathbb{C}$  which is non-degenerate and alternating, i.e.  $\omega(v, v) = 0$  for all  $v \in V$ .

For  $V$  symplectic define

$$\mathrm{Sp}(V) := \{g \in \mathrm{GL}(V) \mid \omega(gv, gw) = \omega(v, w)\} \leq \mathrm{GL}(V) .$$

# Symplectic Groups

Let  $V$  be a finite dimensional  $\mathbb{C}$ -vector space.

The space  $V$  is called **symplectic** if there is a bilinear form  $\omega : V \times V \rightarrow \mathbb{C}$  which is non-degenerate and alternating, i.e.  $\omega(v, v) = 0$  for all  $v \in V$ .

For  $V$  symplectic define

$$\mathrm{Sp}(V) := \{g \in \mathrm{GL}(V) \mid \omega(gv, gw) = \omega(v, w)\} \leq \mathrm{GL}(V) .$$

## Example

Let  $\mathfrak{h}$  be a vector space. Then  $\mathfrak{h} \oplus \mathfrak{h}^*$  is symplectic via

$$\omega((v, f), (w, g)) = g(v) - f(w) .$$

For  $W \leq \mathrm{GL}(\mathfrak{h})$  the induced action on  $\mathfrak{h} \oplus \mathfrak{h}^*$  is symplectic.



# Symplectic Groups

Let  $V$  be a finite dimensional  $\mathbb{C}$ -vector space.

The space  $V$  is called **symplectic** if there is a bilinear form  $\omega : V \times V \rightarrow \mathbb{C}$  which is non-degenerate and alternating, i.e.  $\omega(v, v) = 0$  for all  $v \in V$ .

For  $V$  symplectic define

$$\mathrm{Sp}(V) := \{g \in \mathrm{GL}(V) \mid \omega(gv, gw) = \omega(v, w)\} \leq \mathrm{GL}(V) .$$

## Example

Let  $\mathfrak{h}$  be a vector space. Then  $\mathfrak{h} \oplus \mathfrak{h}^*$  is symplectic via

$$\omega((v, f), (w, g)) = g(v) - f(w) .$$

For  $W \leq \mathrm{GL}(\mathfrak{h})$  the induced action on  $\mathfrak{h} \oplus \mathfrak{h}^*$  is symplectic.

**Fact:**  $\mathrm{Sp}(V) \leq \mathrm{SL}(V)$ .

# Linear Quotients

Let  $G \leq \mathrm{GL}(V)$ ,  $|G| < \infty$ .

# Linear Quotients

Let  $G \leq \mathrm{GL}(V)$ ,  $|G| < \infty$ .

Consider

$$V/G := \mathrm{Spec} \mathbb{C}[V]^G \quad \text{"=" space of orbits .}$$

# Linear Quotients

Let  $G \leq \mathrm{GL}(V)$ ,  $|G| < \infty$ .

Consider

$$V/G := \mathrm{Spec} \mathbb{C}[V]^G \text{ “=” space of orbits .}$$

$C_2$

Let  $C_2 := \langle \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \rangle \leq \mathrm{Sp}_2(\mathbb{C})$ . Then

$$\mathbb{C}[u, v, w] / \langle uv - w^2 \rangle \cong \mathbb{C}[x, y]^{C_2} .$$

# Linear Quotients

Let  $G \leq \mathrm{GL}(V)$ ,  $|G| < \infty$ .


Consider

$$V/G := \mathrm{Spec} \mathbb{C}[V]^G \quad \text{"=" space of orbits}.$$

$C_2$

Let  $C_2 := \langle \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \rangle \leq \mathrm{Sp}_2(\mathbb{C})$ . Then

$$\mathbb{C}[u, v, w] / \langle uv - w^2 \rangle \cong \mathbb{C}[x, y]^{C_2}.$$



# Linear Quotients

Let  $G \leq \mathrm{GL}(V)$ ,  $|G| < \infty$ .

Consider

$$V/G := \mathrm{Spec} \mathbb{C}[V]^G \quad \text{"=" space of orbits}.$$


$C_2$

Let  $C_2 := \langle \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \rangle \leq \mathrm{Sp}_2(\mathbb{C})$ . Then

$$\mathbb{C}[u, v, w] / \langle uv - w^2 \rangle \cong \mathbb{C}[x, y]^{C_2}.$$

**Classical fact**

The variety  $V/G$  is smooth if and only if  $G$  is generated by reflections, i.e.  $g \in \mathrm{GL}(V)$  with  $\mathrm{rk}(g - 1) = 1$ .



# Linear Quotients

Let  $G \leq \mathrm{GL}(V)$ ,  $|G| < \infty$ .

Consider

$$V/G := \mathrm{Spec} \mathbb{C}[V]^G \text{ “=” space of orbits .}$$

$C_2$

Let  $C_2 := \langle \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \rangle \leq \mathrm{Sp}_2(\mathbb{C})$ . Then


$$\mathbb{C}[u, v, w] / \langle uv - w^2 \rangle \cong \mathbb{C}[x, y]^{C_2} .$$

**Classical fact**

The variety  $V/G$  is smooth if and only if  $G$  is generated by reflections, i.e.  $g \in \mathrm{GL}(V)$  with  $\mathrm{rk}(g - 1) = 1$ .

**Corollary**

If  $V$  is symplectic and  $G \leq \mathrm{Sp}(V)$ , then  $V/G$  is singular.



# Symplectic Resolutions

## Resolutions


A **resolution** of  $V/G$  is a smooth variety  $X$  and a proper birational morphism  $X \rightarrow V/G$ .



# Symplectic Resolutions

## Resolutions


A **resolution** of  $V/G$  is a smooth variety  $X$  and a proper birational morphism  $X \rightarrow V/G$ .



# Symplectic Resolutions

## Resolutions

A **resolution** of  $V/G$  is a smooth variety  $X$  and a proper birational morphism  $X \rightarrow V/G$ .




If  $G \leq \mathrm{Sp}(V)$ , then  $V/G$  has a symplectic structure.

# Symplectic Resolutions

## Resolutions

A **resolution** of  $V/G$  is a smooth variety  $X$  and a proper birational morphism  $X \rightarrow V/G$ .



If  $G \leq \mathrm{Sp}(V)$ , then  $V/G$  has a symplectic structure.


## Symplectic resolutions (Beauville, 2000)

A **symplectic resolution** of  $V/G$  is a resolution  $\varphi : X \rightarrow V/G$ , where  $X$  is a symplectic variety and  $\varphi$  is an isomorphism of symplectic varieties over the smooth locus.

# Symplectic Resolutions

## Resolutions

A **resolution** of  $V/G$  is a smooth variety  $X$  and a proper birational morphism  $X \rightarrow V/G$ .



If  $G \leq \mathrm{Sp}(V)$ , then  $V/G$  has a symplectic structure.

## Symplectic resolutions (Beauville, 2000)

A **symplectic resolution** of  $V/G$  is a resolution  $\varphi : X \rightarrow V/G$ , where  $X$  is a symplectic variety and  $\varphi$  is an isomorphism of symplectic varieties over the smooth locus.

In general, those do not exist!

# The Classification Problem

## Classification problem

Classify all  $G \leq \mathrm{Sp}(V)$  for which  $V/G$  admits a (projective) symplectic resolution.

# The Classification Problem

## Classification problem

Classify all  $G \leq \mathrm{Sp}(V)$  for which  $V/G$  admits a (projective) symplectic resolution.

We only need to consider irreducible tuples  $(V, \omega, G)$ .

# The Classification Problem

## Classification problem

Classify all  $G \leq \mathrm{Sp}(V)$  for which  $V/G$  admits a (projective) symplectic resolution.

We only need to consider irreducible tuples  $(V, \omega, G)$ .

$\dim V = 2$

If  $G \leq \mathrm{SL}_2(\mathbb{C})$ , then there is always a symplectic resolution.

→ “Kleinian singularities”

# The Classification Problem

## Classification problem

Classify all  $G \leq \mathrm{Sp}(V)$  for which  $V/G$  admits a (projective) symplectic resolution.

We only need to consider irreducible tuples  $(V, \omega, G)$ .

$\dim V = 2$

If  $G \leq \mathrm{SL}_2(\mathbb{C})$ , then there is always a symplectic resolution.

→ “Kleinian singularities”

## Theorem (Verbitsky, 2000)

If  $V/G$  admits a symplectic resolution, then  $G$  is generated by symplectic reflections, i.e.  $g \in G$  with  $\mathrm{rk}(g - 1) = 2$ .



# The Classification Problem

## Classification problem

Classify all  $G \leq \mathrm{Sp}(V)$  for which  $V/G$  admits a (projective) symplectic resolution.

We only need to consider irreducible tuples  $(V, \omega, G)$ .

$\dim V = 2$

If  $G \leq \mathrm{SL}_2(\mathbb{C})$ , then there is always a symplectic resolution.

→ “Kleinian singularities”

## Theorem (Verbitsky, 2000)


If  $V/G$  admits a symplectic resolution, then  $G$  is generated by symplectic reflections, i.e.  $g \in G$  with  $\mathrm{rk}(g - 1) = 2$ .

## Example

Let  $W \leq \mathrm{GL}(\mathfrak{h})$  be a complex reflection group. Then  $W \leq \mathrm{Sp}(\mathfrak{h} \oplus \mathfrak{h}^*)$  is a symplectic reflection group.


# Symplectic Reflection Groups

Classification by Cohen, 1980




# Symplectic Reflection Groups

Classification by Cohen, 1980



# Symplectic Reflection Groups

Classification by Cohen, 1980



# open cases **DONE!**


**DONE!**

**39**

**6**

# Symplectic Reflection Groups

Classification by Cohen, 1980



1. The Classification Problem

2. The Groups in Question

3. Symplectic Reflection Algebras

4. Conclusion

# Symplectically Primitive, Complex Imprimitive Groups

We consider groups  $G$  which are

- **symplectically primitive**, so there is **no** non-trivial decomposition  $V = V_1 \oplus \cdots \oplus V_k$  into symplectic subspaces such that for any  $g \in G$  and any  $i$  there is  $j$  with  $g(V_i) = V_j$ ;
- **complex imprimitive**, so there exists such a decomposition into (not necessarily symplectic) subspaces.

# Symplectically Primitive, Complex Imprimitive Groups

We consider groups  $G$  which are

- **symplectically primitive**, so there is **no** non-trivial decomposition  $V = V_1 \oplus \cdots \oplus V_k$  into symplectic subspaces such that for any  $g \in G$  and any  $i$  there is  $j$  with  $g(V_i) = V_j$ ;
- **complex imprimitive**, so there exists such a decomposition into (not necessarily symplectic) subspaces.

For these groups we have  $\dim V = 4$  (Cohen, 1980), so we may assume  $V = \mathbb{C}^4$ .



# Symplectically Primitive, Complex Imprimitive Groups

We consider groups  $G$  which are

- **symplectically primitive**, so there is **no** non-trivial decomposition  $V = V_1 \oplus \cdots \oplus V_k$  into symplectic subspaces such that for any  $g \in G$  and any  $i$  there is  $j$  with  $g(V_i) = V_j$ ;
- **complex imprimitive**, so there exists such a decomposition into (not necessarily symplectic) subspaces.

For these groups we have  $\dim V = 4$  (Cohen, 1980), so we may assume  $V = \mathbb{C}^4$ .

Four infinite families of groups  $H \leq \mathrm{GL}_2(\mathbb{C})$ , e.g.  $\mu_{6d}T$ ,  $d \in \mathbb{Z}_{\geq 1}$ , leading to  $E(H) \leq \mathrm{Sp}_4(\mathbb{C})$  generated by

$$h^\vee := \begin{pmatrix} h & 0 \\ 0 & (h^\top)^{-1} \end{pmatrix} \text{ for } h \in H, \text{ and } s := \begin{pmatrix} & & 1 \\ & -1 & \\ 1 & & \end{pmatrix}.$$

# Subgroup Structures

Let  $H_0 \leq H$  be the largest complex reflection subgroup.

## Lemma

The group  $H_0$  is primitive (e.g.  $G_5$  for  $H = \mu_6 T$ ) and  $H_0^\vee$  is a normal subgroup of  $E(H)$ .

# Subgroup Structures

Let  $H_0 \leq H$  be the largest complex reflection subgroup.

## Lemma

The group  $H_0$  is primitive (e.g.  $G_5$  for  $H = \mu_6 T$ ) and  $H_0^\vee$  is a normal subgroup of  $E(H)$ .

## Lemma

Any  $E(H)$  contains a dihedral group  $D_d$  as normal subgroup.

# Subgroup Structures

Let  $H_0 \leq H$  be the largest complex reflection subgroup.

## Lemma

The group  $H_0$  is primitive (e.g.  $G_5$  for  $H = \mu_6 T$ ) and  $H_0^\vee$  is a normal subgroup of  $E(H)$ .

## Lemma

Any  $E(H)$  contains a dihedral group  $D_d$  as normal subgroup.

Write  $\mathcal{S}(G) \subseteq G$  for the subset of symplectic reflections.

## Lemma

We have  $\mathcal{S}(E(H)) = \mathcal{S}(H_0^\vee) \dot{\cup} \mathcal{S}(D_d)$ , stable under  $E(H)$ -conjugacy.

1. The Classification Problem
2. The Groups in Question
3. Symplectic Reflection Algebras
4. Conclusion

# Symplectic Reflection Algebras

Deep link to representation theory:

Consider deformations of  $\mathbb{C}[V] \rtimes G$ , called the **symplectic reflection algebras**  $H_c(V, G)$ .

# Symplectic Reflection Algebras

Deep link to representation theory:

Consider deformations of  $\mathbb{C}[V] \rtimes G$ , called the **symplectic reflection algebras**  $H_c(V, G)$ .

## Technical detail

Here  $c : \mathcal{S}(G) \rightarrow \mathbb{C}$  is a  $G$ -conjugacy invariant function.

# Symplectic Reflection Algebras

Deep link to representation theory:

Consider deformations of  $\mathbb{C}[V] \rtimes G$ , called the **symplectic reflection algebras**  $H_c(V, G)$ .

## Technical detail

Here  $c : \mathcal{S}(G) \rightarrow \mathbb{C}$  is a  $G$ -conjugacy invariant function.

**Etingof–Ginzburg, 2002; Ginzburg–Kaledin, 2004**

If  $V/G$  admits a symplectic resolution, then there exists  $c$  such that  $\dim S = |G|$  for all irreducible  $H_c(V, G)$ -modules  $S$ .



# Symplectic Reflection Algebras

Deep link to representation theory:

Consider deformations of  $\mathbb{C}[V] \rtimes G$ , called the **symplectic reflection algebras**  $H_c(V, G)$ .

## Technical detail

Here  $c : \mathcal{S}(G) \rightarrow \mathbb{C}$  is a  $G$ -conjugacy invariant function.

Etingof–Ginzburg, 2002; Ginzburg–Kaledin, 2004

If  $V/G$  admits a symplectic resolution, then there exists  $c$  such that  $\dim S = |G|$  for all irreducible  $H_c(V, G)$ -modules  $S$ .

## Strategy

Find for all  $c$  an irreducible  $H_c(V, G)$ -module  $S$  with  $\dim S \neq |G|$ .

# Symplectic Reflection Algebras

Deep link to representation theory:

Consider deformations of  $\mathbb{C}[V] \rtimes G$ , called the **symplectic reflection algebras**  $H_c(V, G)$ .

## Technical detail

Here  $c : \mathcal{S}(G) \rightarrow \mathbb{C}$  is a  $G$ -conjugacy invariant function.

## Etingof–Ginzburg, 2002; Ginzburg–Kaledin, 2004

If  $V/G$  admits a symplectic resolution, then there exists  $c$  such that  $\dim S = |G|$  for all irreducible  $H_c(V, G)$ -modules  $S$ .

## Strategy

Find for all  $c$  an irreducible  $H_c(V, G)$ -module  $S$  with  $\dim S \neq |G|$ .

The same strategy was already used for the “improper” symplectic reflection groups.

# Subalgebras

Recall the Lemma:

## Lemma

We have  $\mathcal{S}(E(H)) = \mathcal{S}(H_0^\vee) \dot{\cup} \mathcal{S}(D_d)$ , stable under  $E(H)$ -conjugacy.

# Subalgebras

Recall the Lemma:

## Lemma

We have  $\mathcal{S}(E(H)) = \mathcal{S}(H_0^\vee) \dot{\cup} \mathcal{S}(D_d)$ , stable under  $E(H)$ -conjugacy.

That means we can “split up”  $c$  into two parameters

$$c_1 : \mathcal{S}(H_0^\vee) \rightarrow \mathbb{C} \text{ and } c_2 : \mathcal{S}(D_d) \rightarrow \mathbb{C} .$$

# Subalgebras

Recall the Lemma:

## Lemma

We have  $\mathcal{S}(E(H)) = \mathcal{S}(H_0^\vee) \dot{\cup} \mathcal{S}(D_d)$ , stable under  $E(H)$ -conjugacy.

That means we can “split up”  $c$  into two parameters

$$c_1 : \mathcal{S}(H_0^\vee) \rightarrow \mathbb{C} \text{ and } c_2 : \mathcal{S}(D_d) \rightarrow \mathbb{C} .$$

This gives (sub)algebras

$$H_{c_1}(H_0^\vee) \subseteq H_{c_1}(H^\vee) \subseteq H_{c_1}(E(H)) \longleftrightarrow H_c(E(H)) \longleftrightarrow H_{c_2}(D_d) .$$

# Construction of the Module

$$H_{c_1}(H_0^\vee)$$

$$\cap$$

$$H_{c_1}(H^\vee)$$

$$\cap$$

$$H_{c_1}(E(H))$$

$$\begin{matrix} \uparrow \\ \downarrow \end{matrix}$$

$$H_c(E(H))$$

$$\begin{matrix} \uparrow \\ \downarrow \end{matrix}$$

$$H_{c_2}(D_d)$$

# Construction of the Module

**Remember:** We want to construct an  $H_c(E(H))$ -module of dimension  $\neq |E(H)|$ .

$$H_{c_1}(H_0^\vee)$$

$$\cap$$

$$H_{c_1}(H^\vee)$$

$$\cap$$

$$H_{c_1}(E(H))$$

$$\updownarrow$$

$$H_c(E(H))$$

$$\updownarrow$$

$$H_{c_2}(D_d)$$

# Construction of the Module

**Remember:** We want to construct an  $H_c(E(H))$ -module of dimension  $\neq |E(H)|$ .

**Step 1:** Reduction to  $H_{c_1}(H^\vee)$  and  $D_d$ .

$$H_{c_1}(H_0^\vee)$$

$$\cap$$

$$H_{c_1}(H^\vee)$$

$$\cap$$

$$H_{c_1}(E(H))$$

$$\begin{array}{c} \updownarrow \\ \updownarrow \end{array}$$

$$H_c(E(H))$$

$$\begin{array}{c} \updownarrow \\ \updownarrow \end{array}$$

$$H_{c_2}(D_d)$$



# Construction of the Module

**Remember:** We want to construct an  $H_c(E(H))$ -module of dimension  $\neq |E(H)|$ .

**Step 1:** Reduction to  $H_{c_1}(H^\vee)$  and  $D_d$ .

We can induce any simple  $H_{c_1}(H^\vee)$ -module  $L$  to an  $H_{c_1}(E(H))$ -module  $M$  with  $\dim M = 2 \dim L$ .

$$H_{c_1}(H_0^\vee)$$

$$\cap$$

$$H_{c_1}(H^\vee)$$

$$\cap$$

$$H_{c_1}(E(H))$$

$$\updownarrow$$

$$H_c(E(H))$$

$$\updownarrow$$

$$H_{c_2}(D_d)$$

# Construction of the Module

**Remember:** We want to construct an  $H_c(E(H))$ -module of dimension  $\neq |E(H)|$ .

**Step 1:** Reduction to  $H_{c_1}(H^\vee)$  and  $D_d$ .

We can induce any simple  $H_{c_1}(H^\vee)$ -module  $L$  to an  $H_{c_1}(E(H))$ -module  $M$  with  $\dim M = 2 \dim L$ .

## Theorem

This is an  $H_c(E(H))$ -module if and only if all the constituents of  $M|_{D_d}$  are  $c_2$ -rigid, i.e. if they are isomorphic to a simple  $H_{c_2}(D_d)$ -module.

$$H_{c_1}(H_0^\vee)$$

$$\cap$$

$$H_{c_1}(H^\vee)$$

$$\cap$$

$$H_{c_1}(E(H))$$

$$\Updownarrow$$

$$H_c(E(H))$$

$$\Updownarrow$$

$$H_{c_2}(D_d)$$

# Construction of the Module

**Remember:** We want to construct an  $H_c(E(H))$ -module of dimension  $\neq |E(H)|$ .

**Step 1:** Reduction to  $H_{c_1}(H^\vee)$  and  $D_d$ .

We can induce any simple  $H_{c_1}(H^\vee)$ -module  $L$  to an  $H_{c_1}(E(H))$ -module  $M$  with  $\dim M = 2 \dim L$ .

## Theorem

This is an  $H_c(E(H))$ -module if and only if all the constituents of  $M|_{D_d}$  are  $c_2$ -rigid, i.e. if they are isomorphic to a simple  $H_{c_2}(D_d)$ -module.

**Step 2:** Reduction to  $H_0$ .

$$H_{c_1}(H_0^\vee)$$

$$\cap$$

$$H_{c_1}(H^\vee)$$

$$\cap$$

$$H_{c_1}(E(H))$$

$$\updownarrow$$

$$H_c(E(H))$$

$$\updownarrow$$

$$H_{c_2}(D_d)$$

# Construction of the Module

**Remember:** We want to construct an  $H_c(E(H))$ -module of dimension  $\neq |E(H)|$ .

**Step 1:** Reduction to  $H_{c_1}(H^\vee)$  and  $D_d$ .

We can induce any simple  $H_{c_1}(H^\vee)$ -module  $L$  to an  $H_{c_1}(E(H))$ -module  $M$  with  $\dim M = 2 \dim L$ .

## Theorem

This is an  $H_c(E(H))$ -module if and only if all the constituents of  $M|_{D_d}$  are  $c_2$ -rigid, i.e. if they are isomorphic to a simple  $H_{c_2}(D_d)$ -module.

**Step 2:** Reduction to  $H_0$ .

For  $\lambda \in \text{Irr}(H)$ , we have  $\lambda|_{H_0} \in \text{Irr}(H_0)$  giving rise to a simple  $H_{c_1}(H_0^\vee)$ -module  $L(\lambda|_{H_0})$  with  $\dim L(\lambda|_{H_0}) \leq |H_0|$ .

$$H_{c_1}(H_0^\vee)$$

$$\cap$$

$$H_{c_1}(H^\vee)$$

$$\cap$$

$$H_{c_1}(E(H))$$

$$\Downarrow$$

$$H_c(E(H))$$

$$\Downarrow$$

$$H_{c_2}(D_d)$$

# Construction of the Module

**Remember:** We want to construct an  $H_c(E(H))$ -module of dimension  $\neq |E(H)|$ .

**Step 1:** Reduction to  $H_{c_1}(H^\vee)$  and  $D_d$ .

We can induce any simple  $H_{c_1}(H^\vee)$ -module  $L$  to an  $H_{c_1}(E(H))$ -module  $M$  with  $\dim M = 2 \dim L$ .

## Theorem

This is an  $H_c(E(H))$ -module if and only if all the constituents of  $M|_{D_d}$  are  $c_2$ -rigid, i.e. if they are isomorphic to a simple  $H_{c_2}(D_d)$ -module.

**Step 2:** Reduction to  $H_0$ .

For  $\lambda \in \text{Irr}(H)$ , we have  $\lambda|_{H_0} \in \text{Irr}(H_0)$  giving rise to a simple  $H_{c_1}(H_0^\vee)$ -module  $L(\lambda|_{H_0})$  with  $\dim L(\lambda|_{H_0}) \leq |H_0|$ .

## Lemma

The module  $L(\lambda|_{H_0})$  is a simple  $H_{c_1}(H^\vee)$ -module as well.

$$\begin{array}{c} H_{c_1}(H_0^\vee) \\ \cap \\ H_{c_1}(H^\vee) \\ \cap \\ H_{c_1}(E(H)) \\ \updownarrow \\ H_c(E(H)) \\ \updownarrow \\ H_{c_2}(D_d) \end{array}$$

1. The Classification Problem
2. The Groups in Question
3. Symplectic Reflection Algebras
4. Conclusion

# Results

## Conclusion

We can construct an  $H_c(E(H))$ -module  $M$  with  $\dim M < |E(H)|$ ,  
if we can find  $\lambda$  such that  $L(\lambda|_{H_0})|_{D_d}$  is  $c_2$ -rigid and  $H_0 \not\leq H$ .

# Results

## Conclusion

We can construct an  $H_c(E(H))$ -module  $M$  with  $\dim M < |E(H)|$ , if we can find  $\lambda$  such that  $L(\lambda|_{H_0})|_{D_d}$  is  $c_2$ -rigid and  $H_0 \not\leq H$ .

## Theorem

Such a module exists except in possibly 73 cases.



# Results

## Conclusion

We can construct an  $H_c(E(H))$ -module  $M$  with  $\dim M < |E(H)|$ , if we can find  $\lambda$  such that  $L(\lambda|_{H_0})|_{D_d}$  is  $c_2$ -rigid and  $H_0 \not\leq H$ .

## Theorem

Such a module exists except in possibly 73 cases.

Using the MAGMA-package CHAMP (Thiel, 2013) we can improve those theoretical bounds and obtain sharp bounds in many cases:

# Results

## Conclusion

We can construct an  $H_c(E(H))$ -module  $M$  with  $\dim M < |E(H)|$ , if we can find  $\lambda$  such that  $L(\lambda|_{H_0})|_{D_d}$  is  $c_2$ -rigid and  $H_0 \not\leq H$ .

## Theorem

Such a module exists except in possibly 73 cases.

Using the MAGMA-package CHAMP (Thiel, 2013) we can improve those theoretical bounds and obtain sharp bounds in many cases:

## Refined Theorem

Such a module exists except in possibly 39 cases. In at least 18 of them no such module exists.

# Results

## Conclusion

We can construct an  $H_c(E(H))$ -module  $M$  with  $\dim M < |E(H)|$ , if we can find  $\lambda$  such that  $L(\lambda|_{H_0})|_{D_d}$  is  $c_2$ -rigid and  $H_0 \not\leq H$ .

## Theorem

Such a module exists except in possibly 73 cases.

Using the MAGMA-package CHAMP (Thiel, 2013) we can improve those theoretical bounds and obtain sharp bounds in many cases:

## Refined Theorem

Such a module exists except in possibly 39 cases. In at least 18 of them no such module exists.

## Final result (so far)

Let  $G \leq \mathrm{Sp}(V)$  be a symplectically primitive complex imprimitive symplectic reflection group. Then the corresponding quotient  $V/G$  does not admit a symplectic resolution except in possibly 39 cases.